

PRESCRIBING INFORMATION TESTOGEL®(testosterone) 16.2MG/G, GEL

For full prescribing information, including side effects, precautions and contraindications, please consult the Summary of Product Characteristics (SPC).

Presentation: Transdermal gel in a multi-dose container, one pump actuation delivers 1.25g of gel containing 20.25mg of testosterone. **Indication:** Testosterone replacement therapy for male hypogonadism when testosterone deficiency has been confirmed by clinical features and biochemical tests. **Dosage and administration:** Cutaneous use. The recommended dose is two pump actuations of gel (i.e. 40.5mg of testosterone) applied once daily. The daily dose should not exceed four pump actuations (81mg testosterone) per day. Adjustment of dosage should be achieved by increments of one pump actuation, usually based on measurements of blood testosterone levels and/or clinical response. The gel should be administered by the patient himself, onto clean, dry, healthy skin on the right and left upper arms and shoulders. **Contraindications:** Cases of known or suspected cancer of the prostate or breast, known hypersensitivity to testosterone or to any other constituent of the gel. **Warnings and precautions for use:** Testosterone deficiency should be clearly demonstrated by clinical features and confirmed by 2 separate blood testosterone measurements. Testosterone levels should be monitored at baseline and at regular intervals during treatment. In addition, in patients receiving long-term androgen treatment the following laboratory parameters should be checked regularly: haemoglobin, haematocrit (to detect polycythaemia), liver function tests and lipid profile. Testogel may affect results of laboratory tests of thyroid function. Risk of pre-existing prostatic cancer should be excluded and the prostate gland and breast monitored during Testogel treatment. Androgens may accelerate the progression of sub-clinical prostate cancer and benign prostate hyperplasia. Testogel should be used with caution in cancer patients at risk of hypercalcaemia and associated hypercalciuria due to bone metastases; regular monitoring of blood calcium levels is recommended in these patients. Testogel may cause oedema with or without congestive cardiac failure in patients suffering from severe cardiac, hepatic or renal insufficiency or ischaemic heart disease. If this occurs, treatment must be stopped immediately. Testogel should be used with caution in patients with ischaemic heart disease. Testosterone may cause a rise in blood pressure and should be used with caution in men with hypertension. Testogel should be used with caution in patients with thrombophilia or risk factors for venous thromboembolism (VTE), as there have been post-marketing reports of thrombotic events in these patients during testosterone therapy. In thrombophilic patients, VTE cases have been reported even under anticoagulation treatment, therefore continuing testosterone therapy after first thrombotic event should be carefully evaluated. In case of treatment continuation, further measures should be taken to minimise the individual VTE risk. Spermatogenesis may be suppressed leading to adverse effects on semen parameters. Gynaecomastia occasionally develops and occasionally persists. Irritability, nervousness, weight gain, prolonged or frequent erections may indicate excessive androgen exposure requiring dosage adjustment. Testogel should be used with caution

in patients with epilepsy and migraine. Do not apply to the genital areas as the high alcohol content may cause local irritation. Testogel is flammable until dry. Testogel can be transferred to other persons by close skin to skin contact. There is limited experience regarding safety and efficacy of Testogel in patients over 65 years of age. Testogel is not indicated for use in women or in children under 18 years of age. Testogel is not a treatment for male impotence or sterility. FOR THE FULL LIST OF WARNINGS AND PRECAUTIONS PLEASE CONSULT SECTION 4.4 OF THE FULL SPC. **Interactions:** May increase the activity of oral anticoagulants. Concomitant administration of testosterone and ACTH or corticosteroids may increase the risk of developing oedema. Testogel may cause changes in insulin sensitivity, glucose intolerance, glycaemic control, blood glucose and glycosylated haemoglobin levels. **Pregnancy and lactation:** Pregnant women must avoid any contact with Testogel application sites. This product may have adverse virilising effects on the foetus. **Undesirable effects:** Local skin reactions include: acne, alopecia, dry skin, skin lesions, contact dermatitis, hair colour changes, rash, sweating, hypertrichosis, application site hypersensitivity, application site pruritus. The following commonly ($\geq 1/100$; $< 1/10$) occur with Testogel: emotional symptoms, prostate specific antigen (PSA) increased, increased haematocrit, increased haemoglobin and increased red blood cell count. The following uncommonly ($\geq 1/1000$ to $< 1/100$) occur with Testogel: malignant hypertension, flushing, phlebitis, diarrhoea, abdominal distention, oral pain, gynaecomastia, nipple disorder, testicular pain, increased erection and pitting oedema. Other adverse reaction identified during post-approval use of Testogel: testis disorder, headache, dizziness, paraesthesia, vasodilation (hot flushes), deep vein thrombosis, dyspnoea, polycythaemia, anaemia, musculoskeletal pain, gynaecomastia, testis disorder, prostate enlargement, oligospermia, benign prostate hyperplasia, impaired urination, anxiety, depression, aggression, insomnia, nausea, asthenia, oedema, malaise and weight increase. In case of severe application site reactions, treatment should be reviewed and discontinued if necessary.

NHS Price: £31.11 per 88g pump pack. **Legal category:** POM. **Marketing Authorisation Number:** PL 28397/0007. **Marketing Authorisation Holder:** Besins Healthcare, Avenue Louise, 287, Brussels, Belgium. **Date of preparation of Prescribing Information:** February 2021 TES/2021/016

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

Adverse events should also be reported to Besins Healthcare (UK) Ltd Drug Safety on 0203 862 0920 or Email: pharmacovigilance@besins-healthcare.com